Investigation Report of Fish and Shellfish Sampled in the Ocean Area ^{Tokyo Electric Power Company} Within 20km Radius of Fukushima Daiichi NPS Sampled in FY 2013 Q4

[Survey of fish/shellfish in which found Multi-nuclide besides radioactive Cs]

(Data summarized on June 13, 2014)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)	
			Sr∗90 (app. 29 years)	reference (the sum of Cs-134 and Cs- 137)
Common skete(muscle)	Around 3km Offshore of Ukedo River (T-S3)	March 28, 2014	0.11	590
Schlegel's black rockfish(muscle)	(River (1-55)	March 17, 2014	0.24	312
Sebastes cheni(muscle)	Around 2km Offshore of Kido River (T-S5)	March 17, 2014	0.36	283
Microstomus achne(muscle)	Around 4km Offshore of Kumagawa (T-S8)	February 3, 2014	0.47	258
Microstomus achne(muscle)	Around 4km Offshore of Kumagawa (T-S8)	March 30, 2014	1.2	246

* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

* The General Environmental TECHNOS Co., Ltd performed Sr-90 analysis.

* survey on the whole fish