Nuclide Analysis Results of Radioactive Materials in the Air at the Upper Part of the Reactor Building of Unit 2, Fukushima Daiichi < 1/2 >

Reference

(Data summarized on March 9)

Place of Sampling	Upper part of reactor building of Unit 2 (central western side of blowout pannel)		Upper part of reactor building of Unit 2 (central northern side of blowout pannel)		Upper part of reactor building of Unit 2 (central northan side of blowout pannel)		Density limit by the announcement of Reactor Regulation (Bq/cm3) (Density limit in the air to which radiation workers
Time of Sampling	07-Mar-12 8:45 ~ 10:45		07-Mar-12 8:45 ~ 10:45		07-Mar-12 11:00 ~ 13:00		
Detected Nuclides (Half-life)	density of sample (Bq/cm3)	Scaling Factor (/)	density of sample (Bq/cm3)	Scaling Factor (/)	density of sample (Bq/cm3)	Scaling Factor (/)	breathe in the section 4 of the appendix 2)
I-131 (about 8 days)	ND	•	ND	,	ND	ı	1E-03
Cs-134 (about 2 years)	1.3E-05	0.01	1.1E-05	0.01	1.4E-05	0.01	2E-03
Cs-137 (about 30 years)	1.8E-05	0.01	1.2E-05	0.00	1.9E-05	0.01	3E-03

^{*} The value of radioactivity density is the sum of the value of volatile nuclide's density and the value of particulate nuclide's density.

O.OE - O means O.O x 10-O

Data of other nuclides are under examination.

The followings show the detection limits.

Volatile: I-131: approx. 5E-6Bq/cm3, Cs-134: approx. 7E-6Bq/cm3, Cs-137: approx. 8E-6Bq/cm3

Particulate: I-131: approx. 2E-6Bq/cm3

Please note that these nuclides are sometimes detected even when they are below the limits, contingent on the detector or samples.

^{*} In the case that two or more kinds of nuclides exist, sum of each scaling factor to the density limit is compared with 1.

^{* &}quot;ND" means the sampled data is below measurable limit.

Nuclide Analysis Results of Radioactive Materials in the Air at the Upper Part of the Reactor Building of Unit 2, Fukushima Daiichi < 2/2 >

Reference

(Data summarized on March 9)

Place of Sampling	Upper part of react of Unit 2 (central northern si out pannel)						Density limit by the announcement of Reactor
Time of Sampling	7-Mar-12 11:00 ~ 13:00						Regulation (Bq/cm3) (Density limit in the air to which radiation workers
Detected Nuclides (Half-life)	density of sample (Bq/cm3)	Scaling Factor (/)	density of sample (Bq/cm3)	Scaling Factor (/)	density of sample (Bq/cm3)	Scaling	breathe in the section 4 of the appendix 2)
I-131 (about 8 days)	ND	-					1E-03
Cs-134 (about 2 years)	2.5E-05	0.01					2E-03
Cs-137 (about 30 years)	3.1E-05	0.01					3E-03

^{*} The value of radioactivity density is the sum of the value of volatile nuclide's density and the value of particulate nuclide's density.

O.OE - O means O.O x 10-O

Data of other nuclides are under examination.

The followings show the detection limits.

Volatile: I-131: approx. 3E-6Bq/cm3, Cs-134: approx. 7E-6Bq/cm3, Cs-137: approx. 8E-6Bq/cm3

Particulate: I-131: approx. 2E-6Bq/cm3

Please note that these nuclides are sometimes detected even when they are below the limits, contingent on the detector or samples.

^{*} In the case that two or more kinds of nuclides exist, sum of each scaling factor to the density limit is compared with 1.

^{* &}quot;ND" means the sampled data is below measurable limit.