

Nuclide Analysis Results of Fish <Sampled from the Port Area of the Fukushima Daiichi Nuclear Power Station>

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Black sea bream (muscle)	Port area of Fukushima Daiichi NPS (Near port entrance)	November 4, 2019	ND(3.0)	ND(3.2)	ND
Marbled sole (muscle) No.1	Port area of Fukushima Daiichi NPS (Near port entrance)	November 4, 2019	ND(2.5)	2.3	2.3
Marbled sole (muscle) No.2	Port area of Fukushima Daiichi NPS (Near port entrance)	November 4, 2019	ND(2.5)	21	21
Flathead (muscle)	Port area of Fukushima Daiichi NPS (Near port entrance)	November 4, 2019	ND(3.1)	11	11
Flatfish (muscle)	Port area of Fukushima Daiichi NPS (In front of shallow draft quay)	November 29, 2019	3.7	46	49.7
Sea raven (muscle)	Port area of Fukushima Daiichi NPS (North side of the east breakwater)	November 27, 2019	ND(3.3)	3.1	3.1
Sebastes cheni (muscle)	Port area of Fukushima Daiichi NPS (North side of the east breakwater)	November 27, 2019	54	830	884
Marbled sole (muscle) No.3	Port area of Fukushima Daiichi NPS (South side of the east breakwater)	November 19, 2019	ND(3.2)	32	32

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.